

^aAdditional Bright Futures recommendations include: Periodic vision and hearing tests for ages 13–21; risk assessment for anemia for ages 13–21; and fluoride supplementation if needed for ages 13–16.

^bImmunizations should be administered according to the most recent ACIP recommendations (<https://www.cdc.gov/vaccines/schedules/index.html>).

Abbreviations:

ACIP = Advisory Committee on Immunization Practices; BRCA = breast cancer susceptibility gene; CRC = colorectal cancer; CVD = cardiovascular disease; HIV = human immunodeficiency virus; HCV = hepatitis C virus; HBV = hepatitis B virus; STI = sexually transmitted infection; USPSTF = U.S. Preventive Services Task Force; WPSI = Women's Preventive Services Initiative.

*Criteria for selective screening

1. **Low-dose aspirin to prevent cardiovascular disease and colorectal cancer:** Calculated 10-year risk of a CVD event $\geq 10\%$; not at increased risk for bleeding; have a life expectancy of at least 10 years; and are willing to take low-dose aspirin daily for at least 10 years.
2. **Diabetes screening and management:** Overweight or obese and have additional CVD risk factors (hypertension, dyslipidemia, abnormal blood glucose levels, diabetes). Specific anticipatory guidance for ages 13–17 may be found in the Bright Futures Guidelines.
3. **Folic acid supplementation:** Sexually active and planning or capable of pregnancy.
4. **Healthy diet and physical activity counseling:** Overweight or obese and have additional CVD risk factors (hypertension, dyslipidemia, abnormal blood glucose levels, diabetes). Specific anticipatory guidance for ages 13–17 may be found in the Bright Futures Guidelines.
5. **Lipid screening:** Familial dyslipidemia, risk factors, or high-risk conditions for age 40–70 years; previous gestational diabetes but not previously diagnosed with diabetes mellitus when not pregnant for age ≥ 13 years.
6. **Osteoporosis screening:** 10-year fracture risk equivalent to an average-risk 65-year old woman based on specific risk factors (parental history of hip fracture, smoking, white race, excess alcohol consumption, low body weight).
7. **Statin use to prevent CVD:** Age 40 to 75 years; one or more CVD risk factors (i.e., dyslipidemia, diabetes, hypertension, or smoking); and calculated 10-year risk of a CVD event $\geq 10\%$.
8. **Urinary incontinence screening:** Screen all women age 18 and older and younger women if postpartum.
9. **Gonorrhea and chlamydia screening:** New sex partner, more than one sex partner, a sex partner with concurrent partners, or a sex partner who has an STI; inconsistent condom use among persons who are not in mutually monogamous relationships; previous or coexisting STI; and exchanging sex for money or drugs. Prevalence is also higher among incarcerated populations, military recruits, and patients receiving care at public STI clinics.
10. **Hepatitis B screening:** Born in a country with a prevalence of HBV infection $\geq 2\%$; lack of vaccination in infancy in U.S.-born persons with parents from a country or region with prevalence $\geq 8\%$; HIV-positive persons; injection drug users; and household contacts or sexual partners of persons with HBV infection.
11. **Hepatitis C screening:** One-time screening for asymptomatic adults age 18–79 without known liver disease. Repeat screening and screen at other ages if increased risk: past or current injection drug use; receipt of a blood transfusion before 1992; long-term hemodialysis; born to an HCV-infected mother; incarceration; intranasal drug use; getting an unregulated tattoo; and other percutaneous exposures (e.g., health care workers).
12. **HIV preexposure prophylaxis (PrEP):** Candidates for include 1) heterosexually active women with: a serodiscordant sex partner (i.e., in a sexual relationship with a partner living with HIV); or inconsistent use of condoms during sex with a partner whose HIV status is unknown and who is at high risk; or an STI with syphilis or gonorrhea within the past 6 months; 2) uses injection drugs and shared use of drug injection equipment; or has risk of sexual acquisition of HIV based on above; 3) engaged in transactional sex, such as for money, drugs, or other.
13. **Sexually transmitted infection prevention counseling:** New sex partner, more than one sex partner, a sex partner with concurrent partners, or a sex partner who has an STI; inconsistent condom use among persons who are not in mutually monogamous relationships; previous or coexisting STI; and exchanging sex for money or drugs. Prevalence is also higher among incarcerated populations, military recruits, and patients receiving care at public STI clinics.
14. **Syphilis screening:** Women with HIV; high prevalence communities or populations; history of incarceration; exchanging sex for money or drugs.
15. **Latent tuberculosis infection:** Persons from countries with increased tuberculosis prevalence; living in high-risk congregate settings (e.g., homeless shelters, correctional facilities); exposure to individuals with active tuberculosis, such as health care workers and workers in high-risk congregate settings; immunosuppressed individuals.
16. **Breast cancer screening:** No specific criteria; decisions about screening are made on an individual basis through a shared-decision making process.
17. **Lung cancer screening:** 30 pack-year smoking history and currently smoke or have quit within the past 15 years.
18. **Medications to reduce breast cancer risk:** Major risk factors for breast cancer include increasing age, family history of breast or ovarian cancer (especially among first-degree relatives and onset before age 50 years), history of atypical hyperplasia or other nonmalignant high-risk breast lesions, previous breast biopsy, and extremely dense breast tissue. Models suggest that women with an estimated 5-year breast cancer risk of 3% or greater are likely to have more benefit than harm, although the balance of benefits and harms depends on age, race or ethnicity, the medication used, and whether the patient has a uterus.
19. **Skin cancer counseling:** Fair skin, light hair and eye color, freckles, sunburn easily.
20. **Perinatal depression interventions:** Counseling interventions for women with one or more of the following: a history of depression, current depressive symptoms that may not reach a diagnostic threshold, socioeconomic risk factors such as low income or adolescent or single parenthood, recent intimate partner violence, or mental health-related factors such as elevated anxiety symptoms or a history of significant negative life events.
21. **Preeclampsia prevention with low-dose aspirin:** History of preeclampsia, especially when accompanied by an adverse outcome; multifetal gestation; chronic hypertension; type 1 or 2 diabetes mellitus; renal disease; autoimmune disease (systemic lupus erythematosus, antiphospholipid syndrome).
22. **Diabetes screening after pregnancy:** Previous gestational diabetes but not previously diagnosed with diabetes mellitus when not pregnant.



When women are healthy, communities thrive.



Women's Preventive Services Initiative

Recommendations for Well-Woman Care - A Well-Woman Chart



Health Resources & Services Administration

©2020, Content owned by the ACOG Foundation, all Marks and Rights Reserved. This project was supported by the Health Resources and Services Administration (HRSA) of the U.S. Department of Health and Human Services (HHS) under grant number UHOMC29940, Bright Futures for Women's Health: Standard Practice Guidelines for Well Women Care. This information or content and conclusions are those of the author and should not be construed as the official position nor policy of, nor should any endorsements be inferred by HRSA, HHS, or the U.S. Government.

2020 RECOMMENDATIONS FOR WELL-WOMAN CARE

Preventive care visits provide an excellent opportunity for well-woman care including screening, evaluation of health risks and needs, counseling, and immunizations. *Recommendations for Well-Woman Care – A Well-Woman Chart* was developed by the Women's Preventive Services Initiative (WPSI). The Well-Woman Chart outlines preventive services recommended by the WPSI, U.S. Preventive Services Task Force (USPSTF), and Bright Futures based on age, health status, and risk factors. Additional recommendations for immunizations are provided in a separate table from the Advisory Committee on Immunization Practices. Clinical practice considerations, risk assessment methods, and the age and frequency to deliver services are described in the **Clinical Summary Tables** that accompany the chart.

The Well-Woman Chart provides a framework for incorporating preventive health services for women into clinical practice. These services may be completed at a single visit or as part of a series of visits that take place over time. This information is designed as an educational resource to aid clinicians in providing preventive health services for women, and use of this information is voluntary. This information should not be considered as inclusive of all proper treatments or methods of care or as a statement of the standard of care. It is not intended to substitute for the independent professional judgment of the treating clinician. Variations in practice may be warranted when, in the reasonable judgment of the treating clinician, such course of action is indicated by the condition of the patient, limitations of available resources, or advances in knowledge or technology. While every effort is made to present accurate and reliable information, this publication is provided "as is" without any guarantees or warranties of accuracy, reliability, or otherwise, either express or implied. The Chart and Tables are updated annually. The WPSI website (www.womenspreventivehealth.org) has the most up-to-date version of the Chart and Clinical Summary Tables.

PREVENTION SERVICES	AGE (Years)						
	13-17 ^a	18-21 ^a	22-39	40-49	50-64	65-75	>75
GENERAL HEALTH							
Alcohol use screening & counseling	●	●	●	●	●	●	●
Anxiety screening	●	●	●	●	●	●	●
Aspirin to prevent CVD & CRC ¹					○ 50-59		
Blood pressure screening	●	●	●	●	●	●	●
Contraceptive counseling & methods	●	●	●	●	○		
Depression screening	●	●	●	●	●	●	●
Diabetes screening ²	○	○	○	○	○	○	○
Folic acid supplementation ³	○	●	●	●	○		
Healthy diet & activity counseling ⁴	○	○	○	○	○	○	○
Interpersonal violence screening	●	●	●	●	●	●	●
Lipid screening ⁵	○	●	○	●	●	●	
Obesity screening & counseling	●	●	●	●	●	●	●
Osteoporosis screening ⁶					○	●	●
Fall prevention						●	●
Statin use to prevent CVD ⁷				○	○	○	
Substance use assessment	●	●					
Tobacco screening & counseling	●	●	●	●	●	●	●
Urinary incontinence screening ⁸	○	●	●	●	●	●	●
INFECTIOUS DISEASES							
Gonorrhea & chlamydia screening ⁹	●	●	● ≤24 ○ >24	○	○	○	○
Hepatitis B screening ¹⁰	○	○	○	○	○	○	○
Hepatitis C screening (at least once) ¹¹	○	●	●	●	●	●	● <79
HIV preexposure prophylaxis ¹²	○	○	○	○	○	○	○
HIV risk assessment	●	●	●	●	●	●	●
HIV screening (at least once)	● ≥15	●	●	●	●	●	●
Immunizations ^b	●	●	●	●	●	●	●
STI prevention counseling ¹³	●	●	○	○	○	○	○
Syphilis screening ¹⁴	○	○	○	○	○	○	○
Tuberculosis screening ¹⁵	○	○	○	○	○	○	○
CANCER							
Breast cancer screening ¹⁶				○	●	●	○
Cervical cancer screening		● ≥21	●	●	●	●	● ≤65
Colorectal cancer screening					●	●	
Lung cancer screening ¹⁷					○ 55-80	○	○ 55-80
Medications to reduce breast cancer risk ¹⁸				○	○	○	○
Risk assessment for BRCA 1/2 testing		●	●	●	●	●	●
Skin cancer counseling ¹⁹	○	○	○ ≤24				

Recommendations from the WPSI and the USPSTF for preventive services for pregnant and postpartum women are also provided in the Well-Woman Chart. Comprehensive recommendations for pregnant and postpartum women can be found in [ACOG's practice guidelines](#) and other educational materials.

PREVENTION SERVICES for pregnancy provided in addition to age-based services listed above.		PREVENTION SERVICES for postpartum provided in addition to age-based services listed above.	
PREGNANCY		POSTPARTUM	
Anxiety screening	●	Anxiety screening	●
Bacteriuria screening	●	Breastfeeding counseling, services & supplies	●
Breastfeeding counseling, services & supplies	●	Contraceptive counseling & methods	●
Contraceptive counseling & methods	●	Depression and Perinatal Depression Screening ²⁰	●
Depression and Perinatal Depression Screening ²⁰	●	Diabetes screening after gestational diabetes ²²	○
Folic acid supplementation	●	Folic acid supplementation	●
Gestational diabetes screening	●	Interpersonal violence screening	●
Gonorrhea & chlamydia screening	●	Tobacco screening & counseling	●
Hepatitis B screening	●		
HIV screening (each pregnancy)	●		
Interpersonal violence screening	●		
Preeclampsia prevention with low-dose aspirin ²¹	○		
Preeclampsia screening	●		
Rh(D) blood typing	●		
Syphilis screening	●		
Tobacco screening & counseling	●		



The American College of
Obstetricians and Gynecologists
WOMEN'S HEALTH CARE PHYSICIANS

Copyright © 2020 by the ACOG Foundation, updated April 2020



American College of Physicians
Leading Internal Medicine, Improving Lives



NPWH
NURSE PRACTITIONERS
IN WOMEN'S HEALTH
Caring for Women

MEMBERS OF THE ADVISORY PANEL SUPPORT THE WPSI

KEY:

- Recommended by the USPSTF (A or B rating), WPSI, or Bright Futures
- Recommended for selected groups



Women's Preventive Services Initiative